

Intellectual Property Network for Business

[IPN Home](#) | [Search](#) | [Order](#) | [Shopping Cart](#) | [Login](#) | [Help](#)WO9839293A2: 13-THIA PROSTAGLANDINS
FOR USE IN GLAUCOMA THERAPY[View Images \(33 pages\)](#) | [View Cart](#)[Premium Data 1: PDF \(~3140 KB\)](#) | [TIFF \(~2480 KB\)](#) | [Fax](#) | [More choices...](#)**Inventors:**

ZINKE, Paul, W. , 4129 Willow Way Road, Fort Worth, TX 76133, United States of America
HELLBERG, Mark, R. , 5211 Overridge Drive, Arlington, TX 76017, United States of America

Applicant(s):

ALCON LABORATORIES, INC., 6201 South Freeway, Fort Worth, TX 76134-2099, United States of America

Issued/Filed Dates:

Sept. 11, 1998 / March 6, 1998

Application Number:

WO1998US0004505

IPC Class:

C07C 405/00;

Priority Number(s):

March 7, 1997 **US1997060040051**

Designated Countries:

AU, BR, CA, CN, JP, KR, MX, US, **European patent:** AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE

Abstract:

13-thia prostaglandins are useful in the treatment of glaucoma and ocular hypertension. Also disclosed are ophthalmic, pharmaceutical compositions comprising said prostaglandins.

[\[Show "fr" Abstract\]](#)

Attorney, Agent or Firm:

COPELAND, Barry, L.;

Foreign References:

none

(No patents reference this one)

Alternate Searches[Patent Number](#)[Boolean Text](#)[Advanced Text](#)

Nominate this invention for the Gallery...



SEARCH PATENT FULL TEXT
WITH NATURAL LANGUAGE

1 **Premium Data:** Using the premium data links may invoke a charge to your account.

You can check the [pricelist here](#).



[Legal](#) | [IBM](#) | [FAQ](#) | [Feedback](#) | [Contact Us](#)

Use of 13-thiaprostaglandin derivatives - for the treatment of glaucoma and ocular hypertension.

Drug Activity: Ophthalmological; Hypotensive

Mechanism of Action: Prostaglandin

Compound Name: None Given



Use: For the treatment of glaucoma and ocular hypertension (claimed).

Dosage: 0.01-1000 (0.1-100) μ g/eye topically.

Advantage: Reduced side effects, increased discrimination amongst receptors, improved therapeutic profile.

Biological Data: (Ia) was tested for its intraocular pressure (IOP) lowering effect in cynomolgus monkey eyes where ocular hypertension had been induced. Baseline IOP values were determined prior to treatment and 16 hours after the fourth dose. 3.0 mg of (Ia) caused an 18 \pm 3.0% reduction in IOP, compared to a 5.8 \pm 4.0% reduction achieved under identical conditions using PGF $_2\alpha$. Also presented is data showing that (Ia) causes less conjunctival hyperemia, conjunctival swelling and discharge than PGF $_2\beta$.

Chemistry: The use of 13-thia prostaglandins of formula (I) is claimed for the treatment of glaucoma and ocular hypertension.

R1 = CO $_2$ R, ester, CONR $_4$ R $_5$, CH $_2$ OR $_6$ or CH $_2$ NR $_7$ R $_8$; R = H or a cationic salt thereof; R $_4$, R $_5$ = H or alkyl. R $_6$ = H, acyl or alkyl; R $_7$, R $_8$ = H, acyl, or alkyl provided that if either R $_7$ or R $_8$ = acyl, then the other = H or alkyl; n = 0 or 2; R $_2$, R $_3$ = H, alkyl or acyl; B = H, and OH in either configuration, H and F in either configuration, double bonded O, or OCH $_2$ CH $_2$ O; X = (CH $_2$) $_q$ or (CH $_2$) $_q$ O; q = 1-6; Y = 1-6C alkyl group, or a phenyl ring (optionally substituted); or X-Y = (CH $_2$) $_p$ Y $_1$; p = 0-6; Y $_1$ = further defined aromatic moiety; a = single or double bond.

(I) is e.g. (5Z)-(9S, 11R, 15S)-9,11,15-trihydroxy-16-m-chlorophenoxy-13-thia-17,18,19,20-tetranor-5-prostenoic acid isopropyl ester (Ia) (Example II).

33 pages

Drawings 0/0

Authors: Zinke P W; Hellberg M R
Publication Date: 11 September 1998
Language: English
Priority: 07 March 1997 US-040051

Location: Fort Worth, Tex., USA
Document Number: WO9839293-A2
Filed: 06 March 1998 as U04505
Designated States: Regional: AT BE CH DE DK ES FI
FR GB GR IE IT LU MC NL PT SE National: AU BR
CA CN JP KR MX US

WD-98-011852

PP - Cardiovascular

Page - 44

© 1998 Derwent Information